



Aphrodisiac and male fertility enhancing potentials of *Berlinia grandiflora* (Vahl) Hutch. & Dalziel (Leguminosae) using Wistar rats and *Drosophila melanogaster*

Kolapo O Adeleke^{1*}, Adeola O Adedara², Amos O Abolaji², Jones O Moody³

¹ Department of Pharmacognosy, Madonna University, Nigeria

² Department of Biochemistry, University of Ibadan, Ibadan, Nigeria

³ Department of Pharmacognosy, University of Ibadan, Ibadan, Nigeria

Abstract

Inequitable access to assisted reproductive technologies including *in vitro* fertilization to treat infertility remains a challenge for those living in low and middle-income countries. However, medicinal plants have been employed in traditional medicine as alternative remedies to address the problems of infertility. This study was aimed at investigating the male fertility enhancing activities of *Berlinia grandiflora* stem bark.

The powdered sample of *B. grandiflora* (BG) stem bark was macerated in methanol at room temperature for 72 h, the resulting extract (pilot methanol) was tested for possible phytochemicals and evaluated on Wistar rats for toxicity and male fertility enhancing potentials. Proximate analysis was carried out on the stem bark powder and was also successively macerated into n-hexane, dichloromethane, ethyl acetate, and methanol. The extracts were investigated using Wistar rats (sperm indices and hormonal testosterone) and *Drosophila melanogaster* [survival study, mating latency (ML), copulation duration (CD), fly emergence/% fertility] models. Mesterolone (10 mg/kg) and Sildenafil (2 mg/kg) were used as standards for Wistar rats and *Drosophila* models respectively. All data were analyzed using Graph pad prism 5.02, level of significance was placed at $p \leq 0.05$.

Results of acute toxicity and survival study using Wistar rats and *D. melanogaster* respectively showed that BG stem bark extracts are safe above 5 g/kg dose level. 600 mg/kg bwt. of BG pilot methanol extract had the most significant sperm count and Testosterone level respectively ($112.92 \pm 8.97 \times 10^6/\text{mL}$, $0.51 \pm 0.15 \text{ IU/L}$) when compared to the untreated group ($83.27 \pm 3.51 \times 10^6/\text{mL}$, $0.39 \pm 0.03 \text{ IU/L}$) and Mesterolone ($107 \pm 5.08 \times 10^6/\text{mL}$, $1.13 \pm 0.12 \text{ IU/L}$). There was a significant increase in sperm motility in the group treated with 600 mg/kg pilot methanol extract and it was dose-dependent. 400 mg/kg of successive dichloromethane (DCM) and methanol extracts of *B. grandiflora* stem bark showed a significant reduction in ML ($p \leq 0.01$) when compared to the untreated group (1 ml of 0.1% Tween 80 per kg diet). DCM extract showed highest % fertility increase ($50.9 \pm 8.2\%$) in 400 mg/kg ($p \leq 0.001$) relative to negative control (0.00 ± 0.48). Sildenafil (2 mg/kg) had significant mating latency and copulation duration but a non-significant % increase ($6.10 \pm 2.31\%$) infertility

In this study, DCM extract of *Berlinia grandiflora* stem bark was proven to be an effective male fertility enhancer and this has justified its use as a fertility booster in ethnomedicine.

Keywords: *Berlinia grandiflora*, DCM extract, *Drosophila melanogaster*, Mating latency, Male fertility enhancer

Introduction

Infertility is a disease of the reproductive system defined by the failure to achieve a clinical pregnancy after 12 months or more of regular unprotected sexual intercourse (Frey, 2010) ^[11] and there is no other reason, such as breastfeeding or postpartum amenorrhoea (Singh and Pakhiddey, 2015; WHO, 2018) ^[29, 35]. While assisted reproduction technologies (ART) have been available for more than three decades, with more than 5 million children born worldwide from ART interventions such as *in-vitro* fertilization (IVF), these technologies are still largely unavailable, inaccessible, and unaffordable in many parts of the world (Dhont *et al.*, 2012) ^[8], particularly in low and middle-income countries (LMIC). However, medicinal plants have been employed in traditional medicine as alternative remedies to address the problems of infertility at a much-reduced cost and with great accessibility. *Berlinia grandiflora* (Vahl) Hutch. & Dalziel (Leguminosae) ranges in habit from a shrub just 2 - 5 metres tall, to a spreading tree with a dense, rounded, crown that usually grows up to 20 metres tall, exceptionally to 30 metres (Mackinder and Harris, 2006), the bole can be 10 - 70cm in diameter. The tree produces very fragrant, large, conspicuous flowers. The heartwood is pinkish-brown with purple or dark brown veins and frequent resin canals; it is demarcated from the 10 - 15cm wide band of sapwood (Allen and Allen, 1981) ^[1]. *B. grandiflora* is widely distributed in Guinea, Mali, Nigeria, Central Africa, and the Democratic Republic of Congo (Hutchison and Dalziel, 1963) ^[13]. *B. grandiflora* is the most widely spread of the genus and is known to occur within the protected area network. It is called Apado by the Yoruba people and Ububa by the Igbo ethnic groups. *B. grandiflora* is well known among herbalists in the Nigerian traditional medicine practice

for managing several disease conditions such as skin infections, haemorrhoids and liver complications (Joseph *et al.*, 2012) ^[14], cancer (Mike *et al.*, 2010) ^[19], vermifuge, diarrhea and diabetes (Ode *et al.*, 2013) ^[20]. Folks use the leafy twigs decoction as a febrifuge, cholagogue, purgative and antiemetic, while the leaf decoction is also drunk as a tonic (Mackinder and Harris, 2006). It was reported that the leaf and stem of the plant are used by herbalists in South-Eastern Nigeria, to treat microbial infections. Sap from its bark is used for the treatment of sores and wounds, while bark decoctions are administered for the treatment of haemorrhoids and liver complaints. The bark is also used to ease labour during childbirth and gastrointestinal disorders (Joseph *et al.*, 2012) ^[14]. The analgesic activity of stem bark extract was reported by Asuzu *et al.*, (1993) ^[3], the antihelminthic activity of stem bark and its active principle, betulinic acid (Enwerem *et al.*, 2001) ^[10] have been reported. Olatunji *et al.*, 2018 evaluated the anti-tuberculosis activity of the n-hexane, ethyl-acetate, methanol, and aqueous extracts of *B. grandiflora* leaves against *M. bovis*. The aqueous extract of *B. grandiflora* seed was found to have saturated and monounsaturated fatty acids. The high concentration of fatty acids in this extract when in contact with the human aqueous system can lead to severe diarrhoea (Duru *et al.*, 2014) ^[9].

Materials and Methods

Plant collection, authentication, and preparation

Dried barks and voucher specimen of *B. grandiflora* were collected from Km 7, Iseyin- Ado Awaye road, Oja - Agbe, Iseyin local government area, Oyo State, in July 2016. The sample was authenticated at Forestry Herbarium, Ibadan where a voucher specimen (FHI 113219) was deposited. After collection, the bark was cut into pieces, washed, and air-dried under shade, after which it was ground to powder with a Hammermill using a 5KVA motor. The powdered material was weighed and stored in an air-tight container.

Proximate analysis on *B. grandiflora* stem bark (BG)

One gram each of *B. grandiflora* stem bark powder was weighed into six porcelain crucibles, heated for 2 hours in an oven at 105^oc, cooled in desiccators, and weighed. It was heated again at 105^oc hourly, cooled, and weighed until a constant weight was obtained to determine the water loss (%). The methods adopted for the determination of ash values, and extractive values followed the description given by British Pharmacopoeia (1973) ^[5].

Extraction of *B. grandiflora* stem bark (BG)

Two kilograms (1.0 kg) of the pulverized plant material (*B. grandiflora* stem bark: BG) was macerated in methanol for 72 h. It was filtered and the resulting filtrate was concentrated *in vacuo* at 36^oc to obtain 9.36% w/w. The resulting methanol extract (pilot methanol) was used for acute toxicity study and pilot fertility assay. Another batch of the stem bark powder (2.0 kg) was subjected to successive extraction by maceration in n-hexane, dichloromethane, ethyl acetate, and methanol for 72 hours, respectively. Each filtered extract was separately concentrated *in vacuo* to give n-hexane, dichloromethane, ethyl acetate, and methanol extracts which were also used for bioassay.

Phytochemical analysis of *B. grandiflora* stem bark

All phytochemical procedures were carried out using standard methods as described by Sofowora (1993), and Trease and Evans (1989).

Toxicity study and fertility study of *B. Grandiflora* stem bark pilot methanol extract on male Wistar rats

Nine rats weighing 130 - 150g were used for the determination of acute toxicity (1st phase of Lorke's method (Lorke, 1983).

Another batch of thirty rats was employed for the evaluation of liver and kidney function, and fertility study. The animals were divided into groups of 6 animals each. Groups 1-3 were administered methanol extract of *B. grandiflora* stem bark orally at 200, 400, and 600 mg/kg. Mesterolone (10mg/kg) (Group IV) and 0.1% Tween 80 (1 mL/kg) (Group V) served as positive and negative control respectively. The administration was carried out at 9.00 am daily for 14 days. Tween 80 (0.1%) was used to dissolve the extract to enhance solubility and this necessitated its use as the negative control.

Fertility study of *B. grandiflora* stem bark successive extracts on male Wistar rats

The animals were divided into 14 groups of 6 animals each and administered accordingly for 14 days as follows.

Group 1 received 1 ml/kg of 0.1% Tween 80

Group 2 received 10 mg/kg of Mesterolone

Groups 3-5 received 100, 200, and 400 mg/kg, of *B. grandiflora* stem bark n-hexane extract, respectively.

Groups 6-8 received 100, 200, and 400 mg/kg *B. grandiflora* stem bark DCM extract, respectively.

Groups 9-11 received 100, 200, and 400 mg/kg *B. grandiflora* stem bark ethyl acetate extract, respectively

Groups 12-14 received 100, 200, and 400 mg/kg *B. grandiflora* stem bark methanol extract, respectively.

Blood collection and semen analysis

On the 15th day i.e. a day post final extract administration, blood was collected through the orbital sinus of each of the animals in all the groups. The animals were sacrificed by cervical dislocation. The organs such as testes,

kidneys, liver, and epididymis were harvested and weighed. The blood collected was centrifuged at 3000 rpm for 10 minutes to obtain serum which was tested for Testosterone, alanine aminotransferase (ALT), Aspartate aminotransferase (AST), Alkaline phosphatase (ALP), Blood Urea Nitrogen (BUN) and Serum Creatinine levels (Reitman *et al.*, 1957). Testosterone level was determined at Madonna University Teaching Hospital (MUTH) using ELISA random Kit while the toxicity profile (EIA, England) was assayed at the Chemical Pathology Department, Faculty of Veterinary Medicine, University of Ibadan. The right epididymis was removed and the caput was lacerated on a glass slide using a warm (27°C) sterile lancet to release the semen (Oyeyemi *et al.*, 2000). The semen was examined for sperm motility, count, and abnormalities.

Sperm motility

Drops of normal saline and eosin stain were added to affect the full motility of the spermatozoa. Average gross motility was scored under the microscope x 40 objectives, % life-death ratio and caudal epididymis volume were determined (Aweda *et al.*, 2010) ^[10].

Sperm count

The sperm concentration was examined using Neubauer Haemocytometer. (Aweda *et al.*, 2010; Osuchukwu *et al.*, 2016) ^[10, 24]

Sperm abnormalities

Different types of abnormalities found in the sperm cells were analyzed using Smith *et al.*, (1977) method. Two drops of a vital stain eosin-negrosin were added to the semen sample on the slide mixed and a semen smear was prepared on a new clean glass slide. The slide was then scored for such abnormalities as rudimentary tail, bent tail, curved tail, coil tail, tailless head, and headless tail. The scoring was done under x 100 magnification using an Olympus microscope.

Hormonal assay

Blood was collected through the orbital sinus of the Wistar rats using a capillary tube. The blood was thereafter centrifuged, and the serum was separated and used for hormonal assay (ELISA, England) according to the manufacturer's instructions.

Drosophila collection and preparation

Adult *Drosophila melanogaster* was obtained from the Department of Biochemistry, *Drosophila* laboratory, University of Ibadan, Nigeria in 2018. The flies were transported in a controlled environment of 15 to 25°C to Madonna University where they were bred at the Pharmacognosy laboratory extension, Madonna University. They were fed with freshly prepared Corn agar media using the procedure of Guruprasad *et al.*, 2010 ^[12] with some modifications. After 8.5 days of the life cycle, the eggs on the diet metamorphosized into imago. Newly emerged flies were used for the experiment.

Survival study on successive extracts of *B. grandiflora* stem bark using *Drosophila*

Fourteen groups of thirty-five (35) flies each of both sexes from the population were transferred into treatment vials and used for the study. The groups were categorized as follows:

Group 1-3 received 100, 200, and 400 mg/kg *B. grandiflora* stem bark n-hexane extract, respectively in addition to the diet.

Groups 4-6 received 100, 200, and 400 mg/kg, *B. grandiflora* stem bark DCM extract, respectively in addition to the diet.

Groups 7-9 received 100, 200, and 400 mg/kg *B. grandiflora* stem bark ethyl acetate extract, respectively in addition to the diet.

Groups 10-12 received 100, 200, and 400 mg/kg *B. grandiflora* stem bark methanol extract, respectively in addition to the diet.

Groups 13 – untreated group; received diet only (negative control).

Group 14 received Sildenafil citrate (positive control).

The flies were carefully transferred at intervals of four days into freshly prepared diets containing the corresponding extracts. The same was done on the untreated group only that it was not fed with the extracts. Each group was carried out in replicates of five. The number of dead flies was counted and recorded daily for 28 days. Any death occurrence at day zero was counted as death due to handling while deaths from day one to the end of the study were termed as death due to tested extract(s). The percentage of surviving flies was calculated based on the number of flies (35) at the onset of the study.

Effect of *B. grandiflora* stem bark extracts on mating latency, copulation duration, and fly emergence of *D. melanogaster*

The procedure of Suchira and Shakunthala (2014) ^[32] was followed with some modifications. Both sexes were aged for 3 days during which the following procedures were undertaken. The flies were separated into virgin females and bachelor males before 8-10 hours of their emergence when the neurons for mating would have been

activated. The males were divided into groups of 5 each and starved for 8 hours. They were then fed for 64 hours with different concentrations of *B. grandiflora* stem bark extracts (100, 200, and 400 mg/kg diet) of n-hexane, dichloromethane, ethyl acetate, and methanol extracts. The females were also divided into groups of five each but they were not fed with the extracts. After 64 hours of treatment of the male flies with the extracts of which both sexes of flies must have been 3 days old, a pair of both sexes was introduced into the mating chamber and observed for a maximum of 1 hour to determine mating latency and copulation duration. After copulation, male and female fruit flies were transferred from the mating chamber to fresh diets (without extract) for 24 hours before laying eggs. At 24 hours intervals, they were transferred into new vials containing a freshly prepared diet for five days. All the vials in which the flies have been transferred within five days were left for observation to enable the eggs laid to develop into adult flies. The newly emerged flies were summed up in comparison to the untreated group to determine percentage fertility. Sildenafil citrate (2 mg/kg) was used as the reference standard.

Data analysis

Data were analyzed using one-way analysis of variance (ANOVA) employing GraphPad Prism version 5.02 for Windows (Graph pad software, San Diego California, USA, www.graphpad.com) at $P < 0.05$. Dunnett multiple comparison test at 95% Confidence Interval of difference was considered significant.

Results and Discussions

The percentage moisture content of *B. grandiflora* does not fall in the specified range for a powdered herbal product by British Pharmacopoeia, (1973) ^[5]. This means *B. grandiflora* stem bark may be susceptible to microbial infection. Other physicochemical analyses are within Pharmacopoeia standards as revealed in Table 1. The phytochemicals of *B. grandiflora* are shown in Table 2 and they are similar to the observation of Olatunji *et al.*, (2018) ^[23] in their work on Phytochemical evaluation of the same plant. The presence of phytochemicals in plants account for medicinal uses in folklore remedies for the treatment of various ailments (Odoh *et al.*, 2020) ^[21]. There was no significant difference in body weight of animals treated with *B. grandiflora* stem bark extracts (Tables 3 and 4). The organs' weights were not negatively affected in all the groups treated with *B. grandiflora* stem bark extracts unlike in Mesterolone which elicited a reduction in organ weights as shown in Fig 1. Also, all the biochemical indices were normal, the concentrations of Aspartate aminotransferase, Alanine aminotransferase, Alkaline phosphate, Bilirubin urea nitrogen and Creatinine were not significantly different from the control group indicating that the extracts at the tested doses did not have any hazardous effect on the liver and the kidney (Table 5). These biochemical indices, if altered, will impair the normal functioning of the organs (Appidi *et al.*, 2009) ^[2]. The pilot methanol extract showed a dose dependent increase in percentage sperm motility with significant activity at 600 mg/kg (Fig 2) but livability was not affected and there was no significant difference in sperm motility of animals treated with the successive extracts of *B. grandiflora* (Fig 3). A significant increase in sperm motility is an indication of a beneficial effect on the spermatozoa and will enable easy transport of sperm cells into the ovule (Oladeinde *et al.*, 2007) ^[22]. A significant 36% increase in sperm count was produced by pilot methanol extract (600 mg/kg) compared to the negative control (0.1 % Tween 80) and this was higher than that obtained for the positive (29%) control (10 mg/kg Mesterolone). A significant increase in testosterone level was also observed at 400 and 600 mg/kg of the pilot methanol extract with the highest percentage increase of 31% in 600 mg/kg but was lower than that obtained in 10 mg/kg Mesterolone (190%) as represented in Table 6. The successive extract showed no abnormalities in spermatozoa (Table 7). A significant increase in sperm count was also recorded for DCM 400 mg/kg, 200, and 400 mg/kg of methanol extracts among the groups treated with *B. grandiflora* successive extracts (Table 8). A significant increase in sperm count corresponds to improve fertility (Adeleke *et al.*, 2022) ^[15].

A developmental study is a means of investigating the level of toxicity of the test substance. A fruit fly has an average life span of about 50 days (Taylor *et al.*, 2013) ^[33]. The result of survival study showed no significant difference between the groups treated with different concentrations of the extracts of *B. grandiflora* stem bark (Fig 4) in comparison to the negative control group. This implies that the extracts are not toxic to the flies at the concentrations investigated (Ehigiator *et al.*, 2021) ^[7]. In *Drosophila* species, successful mating depends on male activity and female receptivity, and courtship/mating latency is one of the parameters which indicates the vigor of males. It represents the time between the introduction of male and female flies into the observation chamber and the initiation of mating (Mathew and Krishnamurthy, 2018) ^[18]. A male with high vigor reacts quickly in the presence of a female while a male with less vigor reacts slowly. The mating latency and copulation duration here indicate the aphrodisiac property (Adeleke *et al.*, 2022) ^[15] of *B. Grandiflora* stem bark (Fig 5), it was observed that Sildenafil citrate, 400 mg/kg of DCM, and 400 mg/kg diet of methanol extracts showed a significant reduction in mating latency ($p \leq 0.01$) and prolonged copulation duration but it was not significantly different from the untreated group (0 mg/kg diet). A decrease in mating latency means an increase in the vigor of males (Singh and Singh, 2014) ^[28], the present study showed that DCM and methanol extracts arouse the mating ability of male *Drosophila* flies. The extended duration of copulation allows enough time for the male to introduce more sperms into the female flies (Pankaj *et al.*, 2011) ^[26] as evident in percentage fertility. The percentage fertility of successive extracts of *B. grandiflora* stem bark was also determined. The results obtained revealed a 400 mg/kg diet of DCM and methanol extracts as significantly better concentrations among others. 400 mg/kg of DCM and methanol extracts elicited % fertility increases of $50.9\% \pm 8.2$ and $48.2\% \pm 3.2$ respectively at $p \leq 0.01$ when compared to (negative control) untreated group (0.0 ± 4.8). Sildenafil citrate (2 mg/kg) had a significant

aphrodisiac effect but a non-significant % fertility increase (6.1 ± 8.4) as represented in Fig 7. Increased fly emergence implies that the flies are under the influence of the investigated extracts.

B. grandiflora DCM and methanol stem bark extracts were proven to be effective male fertility enhancers. This study provides scientific data to back up the use of the resins secreted from *B. grandiflora* stem bark in the treatment of male infertility by traditional healers.

Conclusion

The use of *B. grandiflora* stem bark in the management of infertility was justified and since it is relatively non-toxic, it can serve as a lead to the discovery of new drug(s) in the treatment of male infertility. However, further studies are required to isolate the bioactive constituents with male fertility-enhancing agents.

Table 1: Proximate analysis of powdered *B. grandiflora* stem bark

Parameters	Values (%)
Moisture content	15.80±1.20
Total ash	12.00±0.20
Acid insoluble ash	1.00±0.01
Water-soluble ash	5.37±0.04
Alcohol extractive value	8.79±0.80
Water extractive value	4.99±0.23

Table 2: Phytochemical analysis of *B. grandiflora* stem bark

Test	Inference
Saponins	+
Tannins	+
Anthraquinones	-
Terpenoids	+
Alkaloids	+
Flavonoids	+
Cyanogenic glycosides	-
Steroids	+

Key: + = Present, - = Absent

Table 3: Effect of pilot methanol extracts (PME) of *B. grandiflora* stem bark on body weight of Wistar rats per week

Weeks	Distilled water (0.9ml/kg)	10 mg/kg mesterolone	Extract (200mg/kg)	Extract (400mg/kg)	Extract (600mg/kg)
0	141.20±5.46	145.60 ± 3.20	165.20 ± 3.22	173.81± 5.49	167.43± 3.61
1	170.80±3.82	160.40 ± 3.60	178.76 ± 4.33	184.52± 1.37	195.25± 2.51
2	185.40±5.10	156.40 ± 1.76	191.52± 3.28	213.06±3.41	209.58± 4.34

Data are expressed as Mean ± SEM (n=6)

Table 4: Effect of successive extracts of *B. grandiflora* stem bark on body weight of treated rats

Grps/ wks	0	1	2
Dist water	128.4±4.5	146.5±6.3	167.8±5.1
Mest 10 mg/kg	126.6±6.5	141.3±8.3	159.2±4.7
N-H 100	131.1±4.2	147.4±5.9	158.2±3.8
N-H 200	127.2 ± 3.7	141.3 ± 3.2	164.9 ± 4.6
N-H 400	128.4±2.9	141.0±3.9	158.2±6.6
DCM 100	130.4±4.9	153.9±4.1	169.7±7.2
DCM 200	131.4±7.1	147.2±4.7	168.4±3.4
DCM 400	131.0±7.8	142.9±4.8	164.5±5.3
EtOAc 100	129.2±4.8	146.7±8.9	157.5±6.5
EtOAc 200	129.6±4.2	146.2±8.2	154.9±5.6
EtOAc 400	126.4±6.4	143.7±7.3	152.8±8.8
MeOH 100	127.3±4.9	140.1±6.4	143.9±6.2
MeOH 200	127.5±3.1	143.6±2.7	149.3±6.9
MeOH 400	125.7±4.7	138.2±5.3	148.9±8.2

Data expressed as Mean ± SEM (n=6)




Fig 1: Effect of *B.grandiflora* pilot methanol stem bark extract on organ weight of animals after 14 days administration

Table 5: Effects of pilot methanol extracts of *B.grandiflorastem* bark on liver and kidney parameters of the treated animals

Liver/kidney function test (IU/L)	Distilled water	Mesterolone 10 mg/kg	MeOH 200 mg/kg	MeOH 400 mg/kg	MeOH 600 mg/kg
AST	37.7±0.7	40.7±0.3	35.9±0.6	36.7±0.2	38.3±0.8
ALT	28.7±0.3	30.0±0.0	27.8±0.3	27.1±0.6	28±0.9
ALP	116.0±1.0	119.3±0.7	111.0±8.2	116.7±7.3	118±4.7
BUN	11.1±0.1	14.0±0.1	11.5 ±0.1	12.3±0.8	12.8±0.4
CREATININE	0.8±0.0	1.0±0.0	0.7±0.1	0.7±0.2	0.8±0.2

Data expressed as Mean ± SEM (n=6)




Fig 2: *B.grandiflora* pilot methanol stem bark extracts on sperm morphology of animals after 14 days administration. Data expressed as Mean ± SEM (n=6), * = p ≤ 0.05




Fig 3: Effect of *B. grandiflora* stem bark successive extracts on the sperm morphology, N-H: Normal hexane, DCM: Dichloromethane, EtOAc: Ethyl acetate, MeOH: methanol. Data expressed as Mean \pm SEM (n=6).

Table 6: Effect of *B. grandiflora* stem bark pilot methanol extract on sperm indices and testosterone level of treated animals.

Treatment/Parameters	Sperm Count ($10^6/ml$)	Sperm volume (ml)	Testosterone (IU/ml)
Dist water + 0.1ml	83.27 \pm 3.51	0.18 \pm 0.02	0.39 \pm 0.03
Mest 10mg/kg	107.00 \pm 5.08* (29%)	0.16 \pm 0.02	1.13 \pm 0.12*** (190%)
200 mg/kg BG	85.78 \pm 7.10 (3%)	0.18 \pm 0.02	0.38 \pm 0.20 (-3%)
400 mg/kg BG	111.57 \pm 4.32* (34%)	0.16 \pm 0.02	0.45 \pm 0.11* (15%)
600 mg/kg BG	112.92 \pm 8.97* (36%)	0.18 \pm 0.00	0.51 \pm 0.15** (31%)

Data expressed as Mean \pm SEM (n=6), *P \leq 0.05, **P \leq 0.01, *** P \leq 0.001, Mest: Mesterolone

Table 7: Effects of *B. grandiflora* stem bark successive extracts on Total sperm abnormalities. N-H: normal hexane, DCM: dichloromethane, EtOAc: ethyl acetate, MeOH: methanol. Data expressed as Mean \pm SEM (n=6)

Treatment groups	Total sperm abnormalities
Distilled water	402.50 \pm 0.64
Mesterolone	401.25 \pm 0.77
N-H 100 mg/kg	402.68 \pm 0.37
N-H 200 mg/kg	402.41 \pm 0.62
N-H 400 mg/kg	402.86 \pm 0.46
DCM 100 mg/kg	402.57 \pm 0.27
DCM 200 mg/kg	402.68 \pm 0.37
DCM 400 mg/kg	402.41 \pm 0.62
EtOAc 100 mg/kg	402.86 \pm 0.46
EtOAc 200 mg/kg	403.32 \pm 0.52
EtOAc 400 mg/kg	403.43 \pm 0.49
MeOH 100mg/kg	402.41 \pm 0.62
MeOH 200mg/kg	402.86 \pm 0.46
MeOH 400 mg/kg	402.57 \pm 0.27

Table 8: Effect of *B. grandiflora* stem bark successive extracts on Sperm count of Wistar rats. N-H: Normal hexane, DCM: Dichloromethane, EtOAc: Ethyl acetate, MeOH: methanol, Mest: Mesterolone, BG: *B. grandiflora*. Data expressed as Mean \pm SEM (n=6), * = p \leq 0.05

Treatment Groups	Sperm count ($10^6/mL$)	% fertility increase/reduction relative to control
Distilled water (control) ml/kg	97.5 \pm 3.3	-
10mg/kg Mest.	108.0 \pm 3.6	10.8

100 N-H BG	90.3±3.8	-7.4
200 N-H BG	92.3±3.6	-5.3
400 N-H BG	95.0±1.1	-2.6
100 DCM BG	102.2± 4.1	4.8
200 DCM BG	96.0±3.7	-1.5
400 DCM BG	116.3± 6.2*	19.3
100 EtOAc BG	98.0±3.6	0.5
200 EtOAc BG	90.3±2.5	-7.4
400 EtOAc BG	93.0±3.5	-4.6
100 MeOH BG	111.2± 2.5	14.1
200 MeOH BG	120.5± 3.4*	23.6
400 MeOH BG	118.0± 5.4*	21.0




Fig 4: Effect of *B. grandiflora* stem bark successive extracts on 28 days survival study using *Drosophila melanogaster*, n-hex: Normal hexane, DCM: Dichloromethane, EtOAc: ethyl acetate, MeOH: methanol




Fig 5: Effect of *B. grandiflora* stem bark successive extracts on mating latency and copulation duration of *D melanogaster* (fruit flies), N-H = normal hexane, DCM = Dichloromethane, EtOAc = ethyl acetate, MeOH = methanol, a = $P \leq 0.01$, b = $p \leq 0.001$

Table 9: Effect of *B. grandiflora* stem bark extracts on fertility using *D. melanogaster* (fruit flies) emergence.

N-H= n-hexane, DCM = dichloromethane, EtOAc = ethyl acetate, MeOH = methanol, ** = P ≤ 0.01 malenogaster. Behav Genet., 7: 359–372.

Treatment Groups	% Emergence of fly	% Fertility increase/ reduction
Distilled water (0 mg/kg)	100.0 ± 4.8	0.0 ± 4.8
Sildenafil (2 mg/kg)	106.1 ± 2.3	6.1 ± 2.3
N-H 100	84.4 ± 3.6	-15.6 ± 3.6
N-H 200	87.3 ± 4.1	-12.7 ± 4.1
N-H 400	135.7 ± 3.8	35.7 ± 3.8
DCM 100	121.7 ± 3.0	21.7 ± 3.0
DCM 200	148.1 ± 14.4**	48.1 ± 14.4**
DCM 400	150.9 ± 8.2**	50.9 ± 8.2**
EtOAc 100	110.8 ± 6.8	10.8 ± 6.8
EtOAc 200	135.2 ± 14.0	35.2 ± 14.0
EtOAc 400	112.9 ± 1.4	12.9 ± 1.4
MeOH 100	75.1 ± 1.4	-24.9 ± 1.4
MeOH 200	114.5 ± 3.8	14.5 ± 3.8
MeOH 400	148.2 ± 3.2**	48.2 ± 3.2**

Acknowledgment

The authors are using this medium to appreciate the effort of Mr. Aderemi Olaiya for the plant collection and also the assistance of Mr. Adeyemo and Dr. Osiyemi of Forestry Research Institute, Ibadan for the plant identification and authentication. May God rewards you immensely.

References

- Allen ON, Allen NK. The leguminosae: A sourcebook of characteristics, uses, and nodulation. University of Wisconsin, 1981:333:32221-5
- Appidi JR, Yakubu MT, Grierson .DS, Afolayan A.J. Toxicological evaluation of aqueous extracts of *Hermannia incana* Cav. leaves in male Wistar rats. *African Journal of Biotechnology*, 2009;8(10):2016-2020.
- Asuzu, JU, Nwele OC, Anaga AO. The pharmacological activities of the stem bark methanolic extract of *Berlinia grandiflora*. *Fitoterapia*, 1993;64(6):529-534.
- Aweda MA, Usikali RM, Awobajo FO. Effects of exposure to 2.5GHZ microwave radiation on male rats' reproductive system. *International Journal of Current Research*, 2010;2(1):120-128.
- British Pharmacopoeia. Printed in England for Her Majesty's Stationery Office by the University printing house Cambridge, 1973, 66.
- British Pharmacopoeia. *Ash value, Acid insoluble, water-soluble extractive, and Alcohol extractive*. Majesty stationary office London (Appendix xii), 1980:2:1276-1279.
- Ben Enoluomen Ehigiator, Raymond Idujemu Ozolua, Rose Obianuju Egbogu. Evaluation of the sex enhancement and fertility properties of *Waltheria indica* ethanol root extract in *Drosophila melanogaster* (Fruit flies). *International Journal of Pharmacology and Pharmaceutical Research*, 2021;3(1):4-8.
- Dhont N, Busasa R, Gasarabwe A. The double burden of HIV and infertility in Rwanda: what lessons can we learn for infertility care in sub-Saharan Africa? *FVV Obgyn*, Monograph, 2012, 75-80.
- Duru CE, Duru IA, Ikpa CBC, Ibe FC. Chemical and spectra studies of the alleged killer seed of *Berlinia grandiflora*. *Journal of Applied Chemistry*, 2014;7(1):14-18.
- Enwerem NM, Okogun JI, Wambebe CO, Okorie DA, Akah PA. Antihelminthic activity of the stem bark extracts of *Berlinia grandiflora* and one of its principles butulinic acid. *Phytomed*, 2001;8:112-114.
- Frey KA. Male reproductive health and infertility. *Prim Care Clin Office Pract*, 2010;37:643-652.
- Guruprasad BR, Hegde SN, Krishna MS. Seasonal and altitudinal variation of 20 Species of Chamundi hill. *J Insect Sci*, 2010;10:123.
- Hutchison J, Dalziel JM. Flora of West Africa. London University Press, London, 1963:2:74.
- Josephs CG, Ching FP, Nnabuife, AC. Investigation of the antimicrobial potentials of some phytochemical extracts of leaf and stem bark of *Berlinia grandiflora* (Leguminosae) Caesalpinioideae against pathogenic bacteria. *Afr. J. Pharmacol. Ther*, 2012;1(3):92-96.
- Kolapo O Adeleke, Cynthia C Ogbonna, Gideon Alade, and Jones O Moody. Effect of *Croton zambescicus* Muell Arg. (Euphorbiaceae) root extracts on aphrodisiac and fertility using Wistar rats and *Drosophila melanogaster* models. *International Journal of Pharmacognosy and Pharmaceutical Research*, 2022;4(1):1-9.
- Lorke D. A new approach to practical acute toxicity testing; *Arch Toxicol*, 1983;54(4):275-87.
- Mackinder BA, Harris DJ. A Synopsis of the genus *Berlinia* (Leguminosae- Ceasalpinioideae), *Edinburgh J. of Bot.* 63:161-182
- Mathew BB, Krishnamurthy NB. Assessment of Lead Toxicity Using *Drosophila melanogaster* as a Model. *Journal of Clinical Toxicology*, 2018;8:380.

19. Mike O, Soladoye NA, Amusa SO, Raji E, Emmanuel CC, Ayanbamiji AT. Ethnobotanical Survey of Anti-Cancer Plants in Ogun State, Nigeria. *Annals of Biological Research*,2010;1(4):261-273.
20. Ode J, Nwaehujor CO, Nwinyi CF. Evaluation of the Anti-diabetic and Antioxidant Activities of the Methanol Leaf Extract of *Berlinia grandiflora*. *Journal of Medical. Sciences*,2013;13(8):743-748.
21. Odoh Uchenna Estella, Ulasi Ijeoma Nneka. Establishment of Quality Parameters and Pharmacognostical Profiling of *Dialium guinneense* L. (Caesalpinioideae), *World Journal of Innovative Research (WJIR)*,2020;8(5):37-46
22. Oladeinde, FO, Kinyua AM, Laditan AA, Michelin R, Bryant JL, Denaro F *et al*. The effects of *Cnidioscolus aconitifolius* (CA) leaf extract and chlorpropamide on blood glucose and insulin levels in the inbred type 2 diabetic mice, *Cell Molecular Biology*,2007;53(3):34-41
23. Olatunji KT, Aliyu A, Henry OE, Oladosu P. Phytochemical constituents and anti-tuberculosis activity of leaf extracts of *Berlinia grandiflora* (Vahl) Hutch. & Dalziel, Nig J. *Pure and Applied Sciences*,2018;3(2):3250-3256.
24. Osuchukwu IW, Sakpa CL, Ekezie J, Okeke C U, Eke C C, Ezejindu DN. Effect of Leaf extract of *Jatropha tanjorensis* on the testis of Wistar Rats. *Journal of Dental and Medical Sciences*.2016;15(4):66-71.
25. Oyeyemi MO, Akusu MO and Ola-davis OE. Effect of Successive ejaculations on the spermogram of West Africa dwarf goats (*Capra hircus* L.). *Vet Arhiv*,2010;70:215-221.
26. Pankaj P, Guru Prasad BR, Anjaneya Murthy N, Hedge SN. The effect of *Emblica officinalis* diet on lifespan, sexual behaviour, and fitness characters in *Drosophila melanogaster*. *International Journal Research in Ayurveda*,2011;32(2):279-284.
27. Reitman S, Frankel SA. Colorimetric Method for the Determination of Serum Glutamic Oxalacetic and Glutamic Pyruvic Transaminases. *American Journal of Clinical Pathology*,1957;28:56-63.
28. Singh A and Singh B.N. Studies on mating behaviour in the *Drosophila bipectinata* species complex: evidence for sperm displacement. *Current Science*,2014;107(3):511-515.
29. Singh V, Pakhiddey R. Current scenario on genetic basis of infertility. A review. *Acta Medica International*, 2015;2:149-154.
30. Smith KD, Rodriguez-Rigau LJ, Steinberger E. Relationship between indices of semen analysis and pregnancy rate in infertile couples. *Fertile Steril*,1977;28(12):134-9.
31. Sofowora A. Medicinal plants and traditional medicine in Africa. Spectrum Books Ltd, Ibadan, Nigeria, 1993, 289.
32. Suchira PG, Shkunthala V. Effect of different light regimes on fitness of *Drosophila agumbensis* and *Drosophila nagarholensi*, *International Journal of Current Microbiology and Applied Science*,2014;3(10):326-335
33. Taylor ML, Evans JP, Garcia-Gonzalez F. No Evidence for Heritability of Male Mating Latency or Copulation Duration across Social Environments in *Drosophila melanogaster*. *Public Library of Science One*,2013;8(10):12-14.
34. Trease GE, Evans WC. Phytochemical screening: Pharmacognosy, 11th edn, Brailliar Tiridel Can. Macmillian Publishers, London, England, 1989, 235-238.
35. World Health Organization (WHO). International Classification of Diseases, 11th Revision (ICD-11) Geneva: WHO 2018.